SERIAL NO.: 10/602,242 FILED: JUNE 24, 2003

GROUP ART UNIT: 1641

EXAMINER: NELSON C. YANG

PAGE 2

The listing of claims will replace all prior versions, and listings, of claims in the

application:

Listing of Claims:

1. (Previously Amended) A method for detecting and identifying a toxin in a sample,

the method comprises:

providing a dried array comprising a plurality of biological lipid membranes

associated with a surface of a substrate,

wherein the surface comprises a coating of an amine-presenting molecule,

wherein the biological lipid membranes comprise a mixture of a host lipid and a doped

lipid deposited onto the surface;

contacting the array with a solution comprising a toxin; and

monitoring for binding activity of at least one of the biological lipid membranes

with the toxin in the sample.

2. (Previously Amended) The method according to claim 1, wherein the doped lipid

contains a toxin-binding moiety.

3. (Withdrawn) The method according to claim 2, wherein said toxin-binding moiety is

a cell-surface protein.

4. (Previously Amended) The method according to claim 2, wherein the toxin-binding

moiety is a carbohydrate.

5. (Previously Amended) The method according to claim 4, wherein the carbohydrate

moiety is a ganglioside.

SERIAL No.: 10/602242 FILED: JUNE 24, 2003

GROUP ART UNIT: 1641 EXAMINER: NELSON C. YANG

PAGE 3

6. (Withdrawn) The method according to claim 2, wherein the toxin-binding moiety is a

natural lipid, a synthetic lipid, or a lipid composition containing a toxin-binding

receptor, or a purified receptor.

7. (Withdrawn) The method according to claim 6, wherein said toxin-binding moiety is

an ion channel.

8. (Withdrawn) The method according to claim 8, wherein the toxin-binding receptor is

a sodium channel, a potassium channel, a calcium channel, and any combination of ion

channels, an acetylcholine receptor, a ryanodine receptor, a glutamate receptor, a

ceramide, a ganglioside, a cerebroside, sulfatides or cholesterol.

9. (Previously Amended) The method according to claim 1, wherein the biological lipid

membranes are arranged in distinct microspots.

10. (Previously Amended) The method according to claim 1, wherein the toxin has at

least one constituent that is labeled.

11. (Previously Amended) The method according to claim10, wherein the monitoring

step comprises detecting for the presence of the label.

12. (Previously Amended) The method according to claim 1, wherein the monitoring

step comprises detecting directly a physical change due to the binding of the toxin to

the biological lipid membranes.

13. (Previously Amended) The method according to claim 1, wherein the toxin has no

labeled constituent.

SERIAL NO.: 10/602242 FILED: JUNE 24, 2003

GROUP ART UNIT: 1641 EXAMINER: NELSON C. YANG

PAGE 4

14. (Previously Amended) The method according to claim 1, wherein the method

employs a labeled toxin or known compounds with an affinity to the toxin or to a

receptor site of the toxin.

15. (Previously Amended) The method according to claim 1, the toxin detection sample

can be a synthetic or natural toxin, or from a human, animal, plant, food, or

environmental source.

16. (Original) The method of claim 1, wherein the substrate includes a glass, ceramic,

metal-oxide, metal, non-metal, silicon, or polymer material.

17. (Previously Amended) The method according to claim 1, wherein the substrate is

either nano- or micro-porous.

18. (Original) The method according to claim 1, wherein the substrate is configured as a

bead, chip, a slide, a multiwell microplate, or a microcolumn.

19. - 26. (Canceled)

27. (Withdrawn) An array for identifying and detecting a toxin, the array comprising a

plurality of biological membrane probes associated with a surface of a substrate; said

biological membrane containing a toxin-binding moiety.

28. - 41. (Canceled)

SERIAL NO.: 10/602242 FILED: JUNE 24, 2003 GROUP ART UNIT: 1641 EXAMINER: NELSON C. YANG

PAGE 5

42. (Previously Amended) A method for detecting a binding event between a probe and target compound, the method comprising:

providing a dried array comprising a plurality of biological lipid membrane microspots associated with a surface of a substrate, wherein the surface comprises a coating of an amine-presenting molecule, and each of the biological lipid membrane microspots comprises a mixture of a host lipid and a doped lipid deposited onto the surface;

contacting a solution comprising a target compound with the array of probe biological lipid membrane microspots; and

detecting a binding event between at least one or more of the probe biological lipid membrane microspots with one or more constituents of the target compound.

- 43. (Previously Amended) The method of claim 42, wherein at least one of the constituents of the target compound is labeled and the detection step comprises detecting the presence of the label.
- 44. (Previously Amended) The method of claim 43, wherein the detection of the label is carried out by imaging based on fluorescence, phosphorescence, chemiluminescence, or resonance light scattering emanating from the bound target.
- 45. (Original) The method of claim 42, further comprising washing the substrate of unbound target prior to the detection step.
- 46. (Previously Amended) The method of claim 42, wherein the array of biological lipid membrane microspots is incubated with a labeled target compound and an unlabeled target compound, and the binding event between the unlabeled target compound and the probe is determined by measuring a decrease in the signal of the label due to competition between the labeled target and the unlabeled target compound for the probe.

SERIAL NO.: 10/602242 FILED: JUNE 24, 2003

GROUP ART UNIT: 1641

EXAMENER: NELSON C. YANG

PAGE 6

47. (Original) The method of claim 42, wherein the target is unlabeled and the binding

event is determined by a change in physical properties at the interface.

48. (Original) The method of claim 47, wherein the change in physical properties at the

interface is a change in refractive index or electrical impedance.

49. (Previously Amended) A method for identifying and detecting a toxin in a sample,

the method comprising:

providing a dried array comprising a plurality of biological lipid membrane

microspots associated with a surface of a substrate, wherein the surface comprises a

coating of an amine-presenting molecule, and each of the biological lipid membrane

microspots comprises a mixture of a host lipid and a doped lipid deposited onto the

surface;

contacting a sample comprising an unknown toxin with the array of biological

lipid membrane microspots; and

detecting the binding profile of the unknown toxin to at least one or more of the

biological lipid membrane microspots.

50. (Previously Amended) The method of claim 49, wherein the sample is a biofluid

from a specific infectious tissue, a solution from food or environmental sources, or an

aqueous solution comprising chemical toxins collected or concentrated from a

contaminated gaseous medium.

51. (Previously Amended) The method according to claim 1, wherein the amine-

presenting molecule is γ -aminopropylsilane.

ATTORNEY DOCKET NO.: SP02-143 SERIAL NO.: 10/602242 FILED: JUNE 24, 2003 GROUP ART UNIT: 1641 EXAMINER: NELSON C. YANG PAGE 7

- 52. (Previously Amended) The method according to claim 1, wherein the amine-presenting molecule is selected from the group consisting of poly-lysine, polyethyleneimine, and chitosan.
- 53. (Previously Amended) The method according to claim 42, wherein the amine-presenting molecule is γ-aminopropylsilane.
- 54. (Previously Amended) The method according to claim 42, wherein the amine-presenting molecule is selected from the group consisting of poly-lysine, polyethyleneimine, and chitosan.
- 55. (Previously Amended) The method according to claim 49, wherein the amine-presenting molecule is γ-aminopropylsilane.
- 56. (Previously Amended) The method according to claim 49, wherein the amine-presenting molecule is selected from the group consisting of poly-lysine, polyethyleneimine, and chitosan.
- 57. (Previously Amended) A method for detecting a binding event between a receptor in a biological lipid membrane and a target compound, said method comprising:

contacting a solution comprising the target compound with a dried array which comprises a plurality of biological lipid membranes associated with a surface of a substrate, wherein each of the biological lipid membranes comprise a mixture of a host lipid and a receptor of interest and is deposited onto a coating of the surface of the substrate;

detecting a binding event between one or more receptors in the biological lipid membranes and one or more constituents of the target compound, and

wherein the coating of the surface comprises an amine-presenting molecule or a silane.

SERIAL NO.: 10/602242

Filed: June 24, 2003 GROUP ART UNIT: 1641

EXAMINER: NELSON C. YANG

PAGE 8

58. (Previously Amended) The method of claim 57, wherein the coating consists of a

coating of the amine-presenting molecule.

59. (Previously Amended) The method of claim 58, wherein the amine-presenting

molecule is selected from the group consisting of γ-aminopropylsilane, polyamine, and

chitosan.

60. (Previously Amended) The method of claim 57, wherein the coating comprises a

coating of the silane.

61. (Previously Amended) The method of claim 60, wherein the silane comprises a

hydroxyl, a carboxyl, a phosphate, a sulfonated, or a thiol group.